

Molecular Detection of *Campylobacter* Species: Comparison of *16SrRNA* with *slyD*, *cadF*, *rpoA*, and *dnaJ* Sequencing

Elahe Ghorbani marghmaleki¹, Azam Ahmadi*¹,
Mohammad Arjomandzadegan¹, Majid Akbari¹, Aysan Karamghoshchi¹

Abstract

Background: *Campylobacter* spp. are the main cause of human gastroenteritis. The *16SrRNA* sequencing is one of fast molecular method to detect this fastidious. In this study, we compared the sequencing of *16srRNA* genewith four housekeeping genestodetect *Campylobacter* spp. in patients with diarrhea and healthy people.

Methods: 60 samples of *Campylobacter* DNA extracted from stool samples of 30 patients with diarrhea and 30 healthy people were used. In order to detect *Campylobacter*, we designed primers for proliferation of *16SrRNA*, *cadF*, *dnaJ*, *slyD*, and *rpoA* genes using Primer 3, Mega 4.0 and Blast software. Then the PCR products were sequenced using ABI system.

Results: The sequencing showed concordance of PCR-products with deposited sequences in the Gene Bank. Among diarrhea patients, 53.3% of samples were significantly ($p < 0.05$) positive for *slyD* and *cadF* genes and 50% of samples were positive using *16SrRNA*, *rpoA*, and *dnaJ* genes by PCR assay. The average of sensitivity and specificity were found 53.33% and 83.33%, respectively.

Conclusions: Due to various copies of repeated sequences of *16SrRNA* gene, analyzing its amplicons on electrophoresis may be more difficult than the *slyD* and *cadF* genes. According to our results, among the 5 studied genes; the highest detection rate was related to *slyD* and *cadF* genes. Although, *dnaJ* and *rpoA* genes, instead of *16SrRNA* gene, can be considered as appropriate genes for molecular detection of *Campylobacter* bacteria.

Keywords: *cadF*, *Campylobacter*, Diarrhea, Molecular detection, *slyD*.

Introduction

Campylobacter spp. are gram-negative, spiral, and comma-shaped bacteria. *Campylobacter* spp. require microaerophilic conditions (10% CO₂, 5% O₂) and also temperatures 37- 42 °C to grow (1). The size of *Campylobacter* genome is 1.6 to 7.1 MBp (2). Virulence factors of *Campylobacter* such as motility, adhesion, invasion to host cell, and production toxin contribute in pathogenicity and causing host cell necrosis (3, 4). Among virulence genes of *Campylobacter*, *flaA*, *flaB*, *racR*, *dnaJ*, and, *slyD* are known to be responsible for pathogenesis, while *ciaB* and *pldA* contribute in gene expression and colonization. Also, *cdtA*, *cdtB*, and *cdtC* genes are

known to be responsible for production of cytotoxin (5). The *slyD* gene encodes Peptidyl Prolyl cis/ trans-isomerase which has role in the synthesis of amino acids(6). Small subunit ribosomal RNA (*16SrRNA*) gene is one of the most important target genes in molecular studies on the bacterial evolution and epidemiology (7). The *16SrRNA* gene is considered as a gold standard gene for estimating of phylogenetic diversity in microbial communities (8,9); Nevertheless, the widespread usage of *16SrRNA*, it can limit several aspects of the results as its high copy numbers in the genome, 1-15 or more copies (10). The copy numbers can

I: Infectious Diseases Research Center (IDRC), Arak University of Medical Sciences, Arak, Iran.

*Corresponding author: Azam Ahmadi; Tel: +98 34173506; E-mail: ahmadia22@yahoo.com.

Received: 19 May, 2020; Accepted: 8 Jun, 2020.

differs between *Campylobacter* spp. and only a limited number of bacterial genomes have one copy of *16SrRNA* gene that its varies may be simultaneous with increasing number of copies (11). Also, the sequence of *16SrRNA* gene can be different among bacteria communities (12).

One of the main causes of diarrhea in humans is *Campylobacter* infection and its detection using bacterial culture is not easy. Therefore, accurate identification of this gene by molecular methods is very important.

The purpose of this study is to compare the detection of *Campylobacter* using the *16SrRNA* gene with *slyD*, *cadF*, *rpoA*, and *dnaJ* genes for determination in patients with diarrhea admitted to Amirkabir Hospital in Arak and healthy individuals people referred to the Arak health center in 2017-2018.

Materials and methods

Primary isolation

This study was approved by the ethics committee of Arak University of Medical Sciences (Ethics code: IR.ARAKMU.REC.1397.229). We used 60 DNA samples available in DNA Bank of Infectious Diseases Research Center of Arak University of Medical Sciences, Iran. DNA samples were extracted from human stools and included 30 samples of patients with diarrhea and 30 fecal specimens from healthy individuals exposed to poultry meat. Because the chicken is a major host for the transmission of *Campylobacter* species.

Detection of *Campylobacter* genus by Polymerase chain reaction (PCR) and Sequencing

For each sample, five housekeeping genes, *16SrRNA*, *cadF*, *rpoA*, *dnaJ*, and, *slyD* were used in PCR. Properties of primers are shown in Table 1. The reaction mixture in final volume was 15 µl, contained 6.2 µl of Master mix (ID No: 5200300-1250, YTA, Iran), 50 ng genomics DNA, and 0.7 µl (10 pmol) of each primer (Copenhagen, Denmark). The PCR was

performed in Thermocycler device (Eppendorf, Germany) under the following temperature conditions: denaturation at 95 °C for 5 min, 35 cycles including initial denaturation of 95 °C for 1 min, annealing 56.4 °C (*16SrRNA* gene) for 55 sec, Extension 72 °C for 1 min and final Extension 72 °C for 10 min. Annealing temperature for each gene is provided in Table 1. DNA extracted from *Campylobacter* colonies (from previous study Reference?) and distilled water were used as positive and negative control, respectively. The identification of *Campylobacter* at genus level from the extracted DNA of fecal specimen was done based on the size of amplicons on agarose 1% gel (Gene Fanavaran, Iran) in horizontal electrophoresis system (Padide Nozhen Pars, Iran) and gel documentation (Quantum ST4, Germany). Then, PCR products were sent to Macrogen company (South Korean) for sequencing and final confirmation by ABI Applied Biosystems 3730xl instrument.

Statistical Analysis

Statistical analyses were carried out using MedCal 18.11. P values less of 0.05 were considered as statistically significant.

Results

PCR results of housekeeping genes

Among of 60 samples, 21 samples (35%), 21 samples (35%), 20 samples (33.33%), samples (33.33%), 20 samples (33.33%), and 20 samples (33.33%) were positive for *Campylobacter* by *slyD*, *cadF*, *dnaJ*, *rpoA*, and *16SrRNA*, respectively Table 2).

Sequencing

Results of sequencing were analyzed by using Mega4 and Chromas software. Figure 1 (a-e) showed amplifications and sequencing results of different genes in present study. Data were compared with sequences of gene bank and confirmed statistically significant using BLAST software.

Table 1. Properties of primers used to detect of *Campylobacter* genus.

Primers ID	Sequence (5' to 3')	Target gene	Size (bp)	Annealing temperature (°C)
<i>slyD-F</i> <i>slyD-R</i>	GCGAAGGTGAAAATGGCGAA GATGATCGTGTCCATGICCG	<i>slyD</i>	216	62
<i>MD16S-F</i> <i>MD16S-R</i>	ATCTAATGGCTTAACCATTA AAC GGACGGTAACTAGTTTAGTATT	<i>16SrRNA</i>	857	56.4
<i>cadF-F</i> <i>cadF-R</i>	TAAAAGCGGTGGATTTGGAC CAGGACATTTTGCTTGTTGGA	<i>cadF</i>	218	54
<i>rpoA-F</i> <i>rpoA-R</i>	CGAGCTTGCTTTGATGAGTG AGTTCACAGGAAAACCTA	<i>rpoA</i>	121	52
<i>dnaJ-F</i> <i>dnaJ-R</i>	GGCAGGGGACAAGTAGGAA CCCTATTGCCACTTTTGCT	<i>dnaJ</i>	227	52

Table 2. Results of amplifications for each group.

Genes	<i>16SrRNA</i>	<i>slyD</i>	<i>cadF</i>	<i>rpoA</i>	<i>dnaJ</i>
Group	Positive number (Percentage)				
Healthy	5 (16.66%)	5 (16.66%)	5 (16.66%)	5 (16.66%)	5 (16.66%)
Patient	15 (50%)	16 (53%)	16 (53%)	15 (50%)	15 (50%)

Table 3. Sensitivity and specificity of PCR.

Total Sample	Gene Diagnostic	Result	Person	Percentage	Diagnostic analysis	Chi-square analysis	p value	DF																																										
30 samples Patient + 30 samples healthy people	Detection With <i>16SrRNA</i> gene	20 positive	15 Patient 5 healthy	25% 8.3%	Sensitivity: 50.00% (31.297%-68.703%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 75.00% (55.529%-87.816%) Negative Predictive Value: 62.50% (52.967%-71.153%) Disease prevalence: 50.00% (36.806%-63.194%)	7.375	0.006	1																																										
		40 Negative	25 healthy	41.7%					30 samples Patient + 30 samples healthy people	Detection With <i>cadF</i> gene	21 positives	16 Patient 5 healthy	26.7% 8.3%	Sensitivity: 53.33% (34.326%-71.658%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 76.190% (57.341%-88.396%) Negative Predictive Value: 64.103% (54.119%-72.998%) Disease prevalence: 50.00% (36.806%-63.194%)	8.717	0.0032	1	39 Negative	25 healthy	41.7%	30 samples Patient + 30 samples healthy people	Detection With <i>slyD</i> gene	21 positive	16 Patient 5 healthy	26.7% 8.3%	Sensitivity: 53.33% (34.326%-71.658%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 76.190% (57.341%-88.396%) Negative Predictive Value: 64.103% (54.119%-72.998%) Disease prevalence: 50.00% (36.806%-63.194%)	8.717	0.0032	1	39 Negative	25 healthy	41.7%	30 samples Patient + 30 samples healthy people	Detection With <i>rpoA</i> gene	20 positive	15 Patient 5 healthy	25% 8.3%	Sensitivity: 50.00% (31.297%-68.703%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 75.00% (55.529%-87.816%) Negative Predictive Value: 62.50% (52.967%-71.153%) Disease prevalence: 50.00% (36.806%-63.194%)	7.375	0.006	1	40 Negative	25 healthy	41.7%	30 samples Patient + 30 samples healthy people	Detection With <i>dnaJ</i> gene	20 positive	15 Patient 5 healthy	25% 8.3%	Sensitivity: 50.00% (31.297%-68.703%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 75.00% (55.529%-87.816%) Negative Predictive Value: 62.50% (52.967%-71.153%) Disease prevalence: 50.00% (36.806%-63.194%)
30 samples Patient + 30 samples healthy people	Detection With <i>cadF</i> gene	21 positives	16 Patient 5 healthy	26.7% 8.3%	Sensitivity: 53.33% (34.326%-71.658%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 76.190% (57.341%-88.396%) Negative Predictive Value: 64.103% (54.119%-72.998%) Disease prevalence: 50.00% (36.806%-63.194%)	8.717	0.0032	1																																										
		39 Negative	25 healthy	41.7%					30 samples Patient + 30 samples healthy people	Detection With <i>slyD</i> gene	21 positive	16 Patient 5 healthy	26.7% 8.3%	Sensitivity: 53.33% (34.326%-71.658%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 76.190% (57.341%-88.396%) Negative Predictive Value: 64.103% (54.119%-72.998%) Disease prevalence: 50.00% (36.806%-63.194%)	8.717	0.0032	1	39 Negative	25 healthy	41.7%	30 samples Patient + 30 samples healthy people	Detection With <i>rpoA</i> gene	20 positive	15 Patient 5 healthy	25% 8.3%	Sensitivity: 50.00% (31.297%-68.703%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 75.00% (55.529%-87.816%) Negative Predictive Value: 62.50% (52.967%-71.153%) Disease prevalence: 50.00% (36.806%-63.194%)	7.375	0.006	1	40 Negative	25 healthy	41.7%	30 samples Patient + 30 samples healthy people	Detection With <i>dnaJ</i> gene	20 positive	15 Patient 5 healthy	25% 8.3%	Sensitivity: 50.00% (31.297%-68.703%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 75.00% (55.529%-87.816%) Negative Predictive Value: 62.50% (52.967%-71.153%) Disease prevalence: 50.00% (36.806%-63.194%)	7.375	0.006	1	40 Negative	25 healthy	41.7%						
30 samples Patient + 30 samples healthy people	Detection With <i>slyD</i> gene	21 positive	16 Patient 5 healthy	26.7% 8.3%	Sensitivity: 53.33% (34.326%-71.658%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 76.190% (57.341%-88.396%) Negative Predictive Value: 64.103% (54.119%-72.998%) Disease prevalence: 50.00% (36.806%-63.194%)	8.717	0.0032	1																																										
		39 Negative	25 healthy	41.7%					30 samples Patient + 30 samples healthy people	Detection With <i>rpoA</i> gene	20 positive	15 Patient 5 healthy	25% 8.3%	Sensitivity: 50.00% (31.297%-68.703%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 75.00% (55.529%-87.816%) Negative Predictive Value: 62.50% (52.967%-71.153%) Disease prevalence: 50.00% (36.806%-63.194%)	7.375	0.006	1	40 Negative	25 healthy	41.7%	30 samples Patient + 30 samples healthy people	Detection With <i>dnaJ</i> gene	20 positive	15 Patient 5 healthy	25% 8.3%	Sensitivity: 50.00% (31.297%-68.703%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 75.00% (55.529%-87.816%) Negative Predictive Value: 62.50% (52.967%-71.153%) Disease prevalence: 50.00% (36.806%-63.194%)	7.375	0.006	1	40 Negative	25 healthy	41.7%																		
30 samples Patient + 30 samples healthy people	Detection With <i>rpoA</i> gene	20 positive	15 Patient 5 healthy	25% 8.3%	Sensitivity: 50.00% (31.297%-68.703%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 75.00% (55.529%-87.816%) Negative Predictive Value: 62.50% (52.967%-71.153%) Disease prevalence: 50.00% (36.806%-63.194%)	7.375	0.006	1																																										
		40 Negative	25 healthy	41.7%					30 samples Patient + 30 samples healthy people	Detection With <i>dnaJ</i> gene	20 positive	15 Patient 5 healthy	25% 8.3%	Sensitivity: 50.00% (31.297%-68.703%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 75.00% (55.529%-87.816%) Negative Predictive Value: 62.50% (52.967%-71.153%) Disease prevalence: 50.00% (36.806%-63.194%)	7.375	0.006	1	40 Negative	25 healthy	41.7%																														
30 samples Patient + 30 samples healthy people	Detection With <i>dnaJ</i> gene	20 positive	15 Patient 5 healthy	25% 8.3%	Sensitivity: 50.00% (31.297%-68.703%) Specificity: 83.33% (65.279%-94.358%) Positive Predictive Value: 75.00% (55.529%-87.816%) Negative Predictive Value: 62.50% (52.967%-71.153%) Disease prevalence: 50.00% (36.806%-63.194%)	7.375	0.006	1																																										
		40 Negative	25 healthy	41.7%																																														




Fig. 1. (A, B) a-e: Sequencing results in Mega4 and chromes software for *16SrRNA* {max score: 67.1, QC: 40%, total score: 67.1, Percent Identity: 82.54 %, E value: 1e-09}, *dnaJ* {max score: 301, QC: 92%, total score: 301, Percent Identity: 94.09%, E value: 5e-80}, *slyD* {max score: 234, QC: 79%, total score: 270, Percent Identity: 87.13%, E value: 7e-60}, *cadF* {max score: 301, QC: 92 %, total score: 301, Percent Identity: 94.09%, E value: 5e-80} and *rpoA* {max score: 115, QC: 76%, total score: 115, Percent Identity: 97.10%, E value: 4e-2}, (C) a-e: Results of PCR product for *16SrRNA*, *dnaJ*, *slyD*, *cadF* and *rpoA* on 1.3% agarose gels. Product size: 857, 227, 216, 218 and 121bp.

Discussion

The increasing isolation of *Campylobacter* from clinical specimens and healthy people has increased its importance in public health (13). Due to the presence of *Campylobacter* in food products with the animal origin, vegetables, and water, this bacterium has been identified as a food-borne pathogen (14). In addition, *Campylobacter* can cause gastrointestinal diseases in humans (gastroenteritis) and abortions in animals (15). Since molecular methods are faster and more accurate than microbial culture methods, in this study we compared molecular diagnostic assay using different genes to detect *Campylobacter* in human fecal samples.

In previous studies, *16SrRNA* gene was mainly used for molecular detection of *Campylobacter* species. Due to the existence of different types of bacterial genomes in the excreted DNA from the feces as well as repeated nucleotide sequences in the *16SrRNA* gene in most bacteria, the identification of the *Campylobacter* is not sufficiently certain by using only *16SrRNA* gene. Regarding the necessity of accurate detection of *Campylobacter*, this study examined the detection of these bacteria by using *cadF*, *rpoA*, *dnaJ*, and *slyD* genes compared with *16SrRNA* gene. Bang et al. in 2003 proposed *cadF*, *flaA*, and *ceuE* genes for determination of *Campylobacter* by PCR assay (16). Ritz et al. in 2009 suggested usage of *rpoA* gene for genotyping of *Campylobacter* species by Multilocus sequence typing (MLST) and evaluation of each gene by using the real time-PCR assay (6). Konkel ME et al. in 1999 used different genes to detect *Campylobacter*, and used *cadF* gene for reproduction and sequencing of gene, and according to this study, this gene can be very suitable for diagnosis of *Campylobacter* (17). Datta et al. in 2003 determined *Campylobacter* spp. with 11 virulence gene by multiplex-PCR assay. Among 11 primers designed for virulence genes, *dnaJ*, *cadF*, and *cdtB* were present in all samples were positive for *Campylobacter* (18).

In this study, not only *16SrRNA* gene was used but also *rpoA*, *cadF*, *slyD*, and *dnaJ* genes were used as housekeeping genes for molecular detection of *Campylobacter*. Among of 30 samples of patients with diarrhea, 15 samples (50%) were

positive for *Campylobacter* by *16SrRNA* gene which was the same with the results of *rpoA* and *dnaJ* genes, While the results of *slyD* and *cadF* genes in patients with diarrhea were 16/30 (53.33%), therefore, the results of these genes were more sensitivity than *16SrRNA*, *dnaJ*, and *rpoA* genes. Also among of healthy people samples, the results of all genes were consistent with each other, and among 30 healthy people samples, *Campylobacter* were identified in five healthy people samples by studied housekeeping genes. These data showed that *cadF* and *slyD* are probably the most suitable genes for accurate detection for *Campylobacter*. According to similar studies, *cadF* and *cdtB* have been identified as suitable genes for the detection this bacterium, and results of this gene are consistent with other studies. The *dnaJ* gene has been reported to be suitable for the diagnosis for *Campylobacter* and its results are consistent with the *16SrRNA* gene. The *rpoA* gene in this study was able to detect 90% of *Campylobacter* isolates, and this result is consistent with other studies. Here, total DNA was extracted from the stool so it is a mixture of genome of different organisms and can the inconsistency with other studies. As shown in Table 3, the sensitivity and specificity of using *cadF* and *slyD* gene for detection of *Campylobacter* by PCR were 53.33% and 83.33%, respectively. The sensitivity and specificity of using these genes have been reported more than *16SrRNA*, *rpoA*, and *dnaJ* housekeeping genes. Therefore, the amplification of the *slyD* and *cadF* genes with specific primers, can be adequate for identification of *Campylobacter* in both groups, patients with diarrhea and healthy people.

In this study, we showed that *Campylobacter* can be detected from a mixture of different genomes with the highest accuracy using housekeeping genes, including *cadF*, *slyD*, *dnaJ*, and *rpoA*, without using the *16srRNA* gene. Probably, replication of *cadF* and *slyD* genes can be considered as a alternative for *16SrRNA* in diagnosis of *Campylobacter*, however research in a larger statistical society is suggested.

The application of PCR molecular method is a more accurate method for diagnosis of *Campylobacter* bacteria in genomic DNA extracted

from the feces. The diagnostic standard in different laboratories is based on the usage of the *16SrRNA* gene. One of important implications of our study is molecular detection of *campylobacter* within template DNA that has been purified from the stool with mixture of genomes of different organisms. In this study, we showed that *Campylobacter* can be detected from a mixture of different genomes with the highest accuracy using housekeeping genes, including *cadF*, *slyD*, *dnaJ* and *rpoA*, Apart from the *16srRNA* gene. Probably, replication of *cadF* and *slyD* genes can be considered

as an alternative for *16SrRNA* in molecular detection of *Campylobacter*.
Campylobacter species.

Acknowledgment

This study is performed in Arak University of Medical Sciences. We are thankful to the staff of Infectious Diseases Research Center of Arak University of Medical Sciences for her technical assistance.

The authors declare they have no conflict of interests.

References

1. Silva J, Leite D, Fernandes M, Mena C, Gibbs PA, et al. *Campylobacter* spp. as a foodborne pathogen: a review. *Front Microbiol.* 2011;2:200.
2. Nuijten PJ, Bartels C, Bleumink-Pluym NM, Gaastra W, van der Zeijst BA. Size and physical map of the *Campylobacter jejuni* chromosome. *Nucleic Acids Res.* 1990;18(21):6211-6214.
3. Altekruse SF, Stern NJ, Fields PI, Swerdlow DL. *Campylobacter jejuni*—an emerging foodborne pathogen. *Emerg Infect Dis.* 1999;5(1):28-35.
4. Wassenaar TM. Toxin production by *Campylobacter* spp. *Clin Microbiol Rev.* 1997;10(3):466-476.
5. Laprade N, Cloutier M, Lapen DR, Topp E, Wilkes G, Villemur R, et al. Detection of virulence, antibiotic resistance and toxin (VAT) genes in *Campylobacter* species using newly developed multiplex PCR assays. *J Microbiol Methods.* 2016;124:41-7.
6. Ritz M, Garenaux A, Berge M, Federighi M. Determination of *rpoA* as the most suitable internal control to study stress response in *C. jejuni* by RT-qPCR and application to oxidative stress. *J Microbiol Methods.* 2009;76(2):196-200.
7. Hugenholtz P, Goebel BM, Pace NR. Impact of culture-independent studies on the emerging phylogenetic view of bacterial diversity. *J bacteriol.* 1998;180(18):4765-4774.
8. Lane DJ, Pace B, Olsen GJ, Stahl DA, Sogin ML, Pace NR. Rapid determination of 16S ribosomal RNA sequences for phylogenetic analyses. *Proc Natl Acad Sci U S A.* 1985;82(20):6955-6959.
9. Ward DM, Weller R, Bateson MM. *16S rRNA* sequences reveal numerous uncultured microorganisms in a natural community. *Nature.* 1990;345(6270):63-5.
10. Klappenbach JA, Saxman PR, Cole JR, Schmidt TM. rrndb: the ribosomal RNA operon copy number database. *Nucleic Acids Res.* 2001;29(1):181-4.
11. Acinas SG, Marcelino LA, Klepac-Ceraj V, Polz MF. Divergence and redundancy of *16S rRNA* sequences in genomes with multiple rrn operons. *J Bacteriol.* 2004;186(9):2629-35.
12. Case RJ, Boucher Y, Dahllöf I, Holmström C, Doolittle WF, Kjelleberg S. Use of *16S rRNA* and *rpoB* genes as molecular markers for microbial ecology studies. *Appl Environ Microbiol.* 2007;73(1):278-88.
13. Leblanc-Maridor M, Garénaux A, Beaudeau F, Chidaine B, Seegers H, Denis M, et al. Quantification of *Campylobacter* spp. in pig feces by direct real-time PCR with an internal control of extraction and amplification. *J Microbiol Methods.* 2011;85(1):53-61.
14. Kang CR, Bang JH, Cho S-I. *Campylobacter jejuni* Foodborne Infection Associated with Cross-contamination: Outbreak in Seoul in 2017. *Infect Chemother.* 2019;51(1):21-27.
15. Bolton DJ. *Campylobacter* virulence and survival factors. *Food microbiol.* 2015;48:99-108.
16. Bang DD, Scheutz F, Oren Gradel K, Nielsen E, Pedersen K, Engberg J, et al. Detection of seven virulence and toxin genes of *Campylobacter jejuni* and *Campylobacter coli* isolates from different sources and cytolethal distending toxin production

suggest potential diversity of pathogenic properties among isolates. *Genome Letters*. 2003;2:62-72.

17. Konkel ME, Kim BJ, Rivera-Amill V, Garvis SG. Identification of proteins required for the internalization of *Campylobacter jejuni* into cultured mammalian cells. *Adv Exp Med Biol*. 1999;473:215-24.

18. Datta S, Niwa H, Itoh K. Prevalence of 11 pathogenic genes of *Campylobacter jejuni* by PCR in strains isolated from humans, poultry meat and broiler and bovine faeces. *J Med Microbiol*. 2003;52(pt 4):345-348.